



Amendments to the Drawings

- P.28, Fig.25
- P.29, Fig.26

On Fig25 and Fig.26, amendments to the numbers on vertical axes are made as shown below.

2→20 4→40 6→60 8→80 10→100

I have attached a replacement sheet of drawings. The amended drawings are identified in the top margin as "Replacement Sheet."

BEST AVAILABLE COPY



DRAWINGS

1/29

Fig. 1

Inhibition effects of anti-HIV agents of the primary processed matter on the syncytium formation of non-infected cells co-cultured with infected cells




Fig. 1b

Inhibition effects of anti-HIV agents of the present invention on the syncytium formation of non-infected cells co-cultured with infected cells





Fig. 2


Inhibition effects of anti-HIV agents of the primary processed matter on HIV production by PHA-stimulated peripheral blood mononuclear cells that were made to be newly infected.



- black part of Kabanoanatake (natural)
- cultured extract
- hyphae cultured and dried by heating
- △— cultured hyphae
- ▲— cultured filtrate

Fig. 2b



Inhibition effects of anti-HIV agents of the present invention on HIV production by PHA-stimulated peripheral blood mononuclear cells that were made to be newly infected.



- black part of Kabanoanatake (natural)
- cultured extract
- hyphae cultured and dried by heating
- △— cultured hyphae
- ▲— cultured filtrate

Fig. 3



Number of viable cells



6/29

Fig. 3b

Number of viable cells







7/29

Fig. 4

ELISA test for HIV P24 antigen yield






8/29

Fig. 4b

ELISA test for HIV P24 antigen yield






9/29


Fig. 5

Anti-HIV effects of pretreated PHA-stimulated peripheral blood mononuclear cells with Kabanoanatake

A The effects of pretreatment HIV with Kabanoanatake



B The effects of target cell pretreatment with Kabanoanatake




* The anti-HIV agents were prepared in PBS solution at the concentration of 3.5 mg/ml.



10/29

Fig. 6

A The effects of pretreatment of target cells with Kabanoanatake



B The effects of addition of Kabanoanatake in various incubation times after target cells pretreatment with anti-HIV agents for approximately one hour






Fig. 6

A-2 The effects of pretreatment of target cells with Kabanoanatake



B-2 The effects of addition of Kabanoanatake in various incubation times after target cells pretreatment with anti-HIV agents for approximately one hour






12/29

Fig. 7


Inhibition effects of anti-HIV agents of the primary processed matter on the syncytium formation of non-infected cells co-cultured with infected cells



* The anti-HIV agents were prepared at the concentration of 3.56 mg/ml.

Fig. 7b

Inhibition effects of anti-HIV agents of the present invention on the syncytium formation of non-infected cells co-cultured with infected cells



* The anti-HIV agents were prepared at the concentration of 3.56 mg/ml.

Fig.8





Fig.8b




Fig. 9

Inhibition effects of various Kabanoanatake of the primary processed matter on the syncytium formation of non-infected cells co-cultured with infected cells







Fig. 9b

Inhibition effects of various Kabanoanatake of the present invention on the syncytium formation of non-infected cells co-cultured with infected cells





18/29

Fig. 10

Report of separation of HIV

July, 18th, 1995

Day of receipt of samples: June, 14th, 1995

(1) Tissue culture infectious dose (TCID)

Total TCID (/ ml)	0
Cell TCID ($/1 \times 10^6$)	0
Plasma TCID (/ ml)	0
Cytopathic effect	0

(2) Anti-HIV antibody in plasma by western blotting methods.

gp160 (env)	gp120 (env)	p65 (pol)	p55 (gag)	p51 (pol)	gp41-43 (env)	p32 (pol)	p24 (gag)	p18 (gag)	p15 (gag)
++	++	++	++	++	++	++	++	++	++

(3) Host range index

(Correspondence column) The virus was not isolated.

(Annotation) Also, in a blood test after three months for the same patient, TCID value was excellent (zero).



19/29

Fig. 10b

Report of separation of HIV

August, 1st, 1998

Day of receipt of samples: June, 14th, 1995

(1) Tissue culture infectious dose (TCID)

Total TCID (/ ml)	0
Cell TCID ($/1 \times 10^6$)	0
Plasma TCID (/ ml)	0
Cytopathic effect	


(2) Anti-HIV antibody in plasma by western blotting methods.

gp160 (env)	gp120 (env)	p65 (pol)	p55 (gag)	p51 (pol)	gp41-43 (env)	p32 (pol)	p24 (gag)	p18 (gag)	p15 (gag)
++	++	++	++	++	++	++	++	++	++

(3) Host range index

(Correspondence column) The virus was not isolated.

Fig.11 Perfect inhibition effects on HIV, in a liquid culture of Kabanoanatake hyphae, AIWro-4, added lignin, for 62 days, under extreme conditions



* A well number below 1 indicates that perfect inhibition effects on HIV is not obtainable.

Fig.12 Cell damage in a liquid culture of Kabanoanatake hyphae, AIW ro-4, when lignin was added






Fig.13 Perfect inhibition effects on HIV in a long-term culture medium of Kabanoanatake hyphae, AIW-27, AIW-4, and lignin, under extreme conditions of restricting the infiltration of oxygen (on the 6th day of the test)



*Diurnal culture temperature was 33°C and night culture temperature was 8°C to 10 °C.
Shaking time was limited to 11 hours per 24 hours.

Fig.14 Cell damage in a liquid culture of Kabanoanatake hyphae, AIW-4 and AIW-27, when lignin was added



*Diurnal culture temperature was 33°C and night culture temperature was 8°C to 10°C.
Shaking time was limited to 11 hours per 24 hours.

Fig.15 Perfect inhibition effects on HIV in a liquid culture of Kabanoanatake hyphae, A-2W- 3 for 34 days, with lignin, under extreme conditions






Fig.16 Cell damage in a liquid culture of Kabanoanatake hyphae, A-2W-3, for 34 days, in the area with lignin, under extreme conditions




* The lines of ①, ② and ③ are the same values, so they are overlapped.

Fig.17 Change in black color tone(500 nm)in a long-term culture test of Kabanoanatake, restricting the infiltration of oxygen




*The control groups (0 g lignin area) for AIW- 4 and AIW-27 were excluded because of growth cessation

Fig. 18 Change in humic acid in a culture medium of Kabanoanatake, under extreme conditions of restricting the infiltration of oxygen




*The control groups (0 g lignin area) for AIW-4 and AIW-27 were excluded because of growth cessation.

Fig.19 Correlation between the amount of lignin-tannin and days of cultures in a long-term culture of Kabanoanatake with lignin, restricting the infiltration of oxygen



*The control groups (0 g lignin area) for AIW-4 and AIW-27 were excluded because of growth cessation

Fig.20 Change in protein amount in a long-term liquid culture test of Kabanoanatake, with lignin, under extreme conditions of restricting the infiltration of oxygen



*The control groups (0 g lignin area) for AIW-4 and AIW-27 were excluded because of growth cessation

Fig.21 Perfect inhibition activity (cells) on HIV, on the 110th day of a liquid culture of Kabanoanatake hyphae, A to E, at the ideal temperature for culture of 25°C




Fig.22 The values of perfect HIV inhibition activity (100%) on the 110th day of a liquid culture of Kabanoanatake hyphae, A to E, at the ideal temperature for culture of 25°C




Fig.23 Change in protein content in a liquid culture of hyphae, AIW-4, with lignin substances (lignosulfonic acid sodium salt acetate and lignosulfonic acid sodium salt)





Fig. 24 Change in protein content in a liquid culture of Kabanoanatake hyphae, A2W-3 and 58-1, when a lignin substance was added






"Replacement Sheet"

28/29

Fig. 25

Incident rate of papillomas, using the 2-stage carcinogenesis model with mouse skin
(average number per mouse)






"Replacement Sheet"

29/29

Fig. 26

The carcinogenetic promotion suppression effects of Compound 1, using the 2-stage carcinogenesis model with mouse skin (percentage)



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.